

M. W. Kalinowski

(Higher Vocational State School in Chełm, Poland)

**On some Cosmological Consequences
of the Nonsymmetric
Kaluza–Klein (Jordan–Thiry) Theory**

Abstract. In the paper we consider some consequences of the Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory. We calculate primordial fluctuation spectrum functions, spectral indices, and first derivatives of spectral indices.

In this paper we develop some ideas from Ref. [1]. We will refer to Ref. [1] in all the details of our considerations.

Let us consider Eq. (3.313) from Ref. [1].

$$\Phi(t) = \frac{(5+3a)\sqrt{5+4a}}{8\alpha_s(A_1 \operatorname{ns}^2(u) + A_2)} + \frac{(1+\sqrt{5+4a})}{2\alpha_s} \quad (1)$$

where

$$u = \sqrt[4]{\frac{Ae^{n\Psi_1}(5Ae^{n\Psi_1} - 3\alpha_s^2 r)}{r^2}} \cdot \sqrt{\sin \frac{\varphi + \pi}{3}} (t - t_0) \quad (2)$$

$$A_1 = \sqrt{\frac{5}{3} - a^2} \sin \frac{\varphi + \pi}{3} \quad (3)$$

$$A_2 = \sqrt{\frac{5}{3} - a^2} \cos \frac{\varphi + 4\pi}{3} + \frac{1}{12}(5 + 6a) \quad (4)$$

and

$$\cos \varphi = -\frac{\sqrt{A}e^{\frac{n}{2}\Psi_1}(9\alpha_s^2 Br^2 + 40Ae^{n\Psi_1})}{12(5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)^{3/2}} \quad (5)$$

$$0 < a = \sqrt{\frac{\alpha_s^2 Br^2}{Ae^{n\Psi_1}}} < 0.3115, \quad (6)$$

ns is the Jacobi elliptic function for the modulus

$$m^2 = \frac{\sin\left(\frac{\varphi}{3}\right)}{\sin\left(\frac{\varphi+2\pi}{3}\right)}, \quad (*)$$

B is an integration constant, r is a radius of manifold of vacuum states (a scale of energy), α_s a coupling constant. A is given by the equation (3.164) and is fixed by

the details of the theory (see Ref. [2]). Ψ_1 is a critical value of a scalar field Ψ (see Ref. [1]). n is a dimension of the group H (see Ref. [1]). Eq. (1) gives a solution to the equation of an evolution of the Higgs field under some special assumptions (in the cosmological background). Due to this solution we are able to calculate $P_R(K)$ —a spectrum of primordial fluctuations in the Universe (see Ref. [3]).

According to the Eq. (3.29)

$$P_R(K) = \left(\frac{H_0}{2\pi} \right)^2 \sum_{\vec{m}, d} \left(\frac{H_0}{\frac{d}{dt}(\Phi_{\vec{m}}^d)} \right)^2 \Big|_{t=t^*}. \quad (7)$$

Using Eq. (1) and the assumption

$$\Phi_{\vec{m}}^d(t) = \delta_{\vec{m}}^d \Phi(t) \quad (8)$$

one gets for $t = t^* = \frac{1}{H_0} \ln \frac{K}{R_0 H_0}$

$$P_R(K) = P_0 \frac{(A_1 + A_2 \operatorname{sn}^2(v))^2 \operatorname{sc}^2(v)}{\operatorname{dn}^2(v)} \quad (9)$$

where H_0 is a Hubble constant

$$P_0 = \frac{12H_0^4 \alpha_s^2 n_1 A^{3/2} e^{(3/2)n\Psi_1} \cdot \sin^{-3} \left(\frac{\varphi + \pi}{3} \right)}{\pi^2 \left(5\sqrt{Ae^{n\Psi_1}} + 3\alpha_s \sqrt{Br^2} \right) \left(5A\sqrt{Ae^{n\Psi_1}} + 4\alpha_s \sqrt{Br^2} \right) (5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)} \quad (10)$$

$$v = (\ln K - t_0 H_0 - \ln(H_0 R_0)) \frac{1}{H_0} \sqrt[4]{\frac{Ae^{n\Psi_1} (5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)}{r^4}} \sqrt{\sin \frac{\varphi + \pi}{3}}, \quad (11)$$

n_1 means a dimension of the manifold $M = G/G_0$ (a vacuum states manifold), $K^2 = \vec{K}^2$ means as usual the square of the length of a fluctuations wave vector. We calculate a spectral index of a power spectrum

$$n_s(K) - 1 = \frac{d \ln P_R(K)}{d \ln K}, \quad (12)$$

$$n_s(K) - 1 = 2C \left(\frac{\operatorname{dn}(v)}{\operatorname{cn}(v) \operatorname{sn}(v)} - m^2 \frac{\operatorname{sn}(v) \operatorname{cn}(v)}{\operatorname{dn}(v)} + \frac{2A_2 \operatorname{sn}(v) \operatorname{cn}(v) \operatorname{dn}(v)}{A_1 + A_2 \operatorname{sn}^2(v)} \right) \quad (13)$$

and an important parameter

$$\begin{aligned} \frac{dn_s(K)}{d \ln K} = 2C^2 & \left[(\operatorname{sn}^2(v) \operatorname{dn}(v) - \operatorname{cn}^2(v) \operatorname{dn}(v) - m^2 \operatorname{sn}(v) \operatorname{cn}(v)) \right. \\ & \times \left(\frac{1}{\operatorname{cn}(v) \operatorname{sn}(v)} + \frac{m^2}{\operatorname{dn}(v)} \right) \\ & + 2A_2 \frac{\operatorname{cn}^2(v) \operatorname{dn}^2(v) - \operatorname{sn}^2(v) \operatorname{dn}^2(v) - m^2 \operatorname{sn}^2(v) \operatorname{cn}^2(v)}{A_1 + A_2 \operatorname{sn}^2(v)} \\ & \left. + 4A_2^2 \frac{\operatorname{sn}^2(v) \operatorname{cn}^2(v) \operatorname{dn}^2(v)}{(A_1 + A_2 \operatorname{sn}^2(v))^2} \right] \end{aligned} \quad (14)$$

where v is given by the formula (11) and all the Jacobi elliptic functions in the formulae (9), (13), (14) have the same modulus

$$m^2 = \frac{\sin \frac{\varphi}{3}}{\sin \frac{\varphi+2\pi}{3}},$$

$$C = \frac{1}{H_0} \sqrt[4]{\frac{Ae^{n\Psi_1} (5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)}{r^4}} \sqrt{\sin \frac{\varphi + \pi}{3}}. \quad (15)$$

In this theory we have some arbitrary parameters which can be translated to parameters A_1, A_2, C and m . Thus we can consider Eqs (9), (13), (14) as parametrized by P_0, A_1, A_2, C and m . Simultaneously v can be rewritten as

$$v = C (\ln K - K'_0) \quad (16)$$

where K'_0 is an arbitrary parameter. Let us remind that in our theory we have arbitrary parameters: r, ζ, ξ, α_s (they could be fixed by some data from elementary particle physics, however, they are free as some parameters of the theory) and integration constants B and t_0 (R_0 also in some sense). The interesting idea for future research consists in applying CMBFAST software [4] to calculate all the cosmological CMB characteristics with mentioned parameters considered as free. Afterwards we should fit them to existing cosmological data [5] as good as possible in order to find some constraints for parameters from our theory.

Let us consider the function $P_R(K)$ from Eq. (9). It is easy to see that the function

$$f(v) = \frac{(A_1 + A_2 \operatorname{sn}^2(v))^2 \operatorname{sc}^2(v)}{\operatorname{dn}^2(v)} \quad (17)$$

is a periodic function

$$f(v) = f(v + 2K(m^2)l), \quad l = 0, \pm 1, \pm 2, \dots, \quad (18)$$

$$K(m^2) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - m^2 \sin^2 \theta}}. \quad (19)$$

Thus we get

$$P_R(K) = P_R(Ke^{l\eta}), \quad (20)$$

$$\eta = \frac{2K(m^2)H_0 \sqrt[4]{\frac{r^4}{Ae^{n\Psi_1}(5Ae^{n\Psi_1} - 3\alpha_s^2 Br^2)}}}{\sqrt{\sin \frac{\varphi + \pi}{3}}} \quad (21)$$

or

$$\eta = \frac{2K(m^2)}{C}. \quad (22)$$

Thus it is enough to consider $P_R(K)$ in the interval

$$e^{-\eta/2} \leq K \leq e^{+\eta/2} \quad (23)$$

(a little similar to the first Brillouin zone). In order to have an interesting cosmological region we should have

$$\eta \simeq 20 \quad (24)$$

which gives us a constraint

$$\frac{2K(m^2)}{C} = \frac{2K(m^2)H_0}{\sqrt{\sin \frac{\varphi+\pi}{3}}} \sqrt[4]{\frac{r^4}{Ae^{n\psi_1}(5Ae^{n\psi_1} - 3\alpha_s^2 Br^2)}} \approx 10. \quad (25)$$

Moreover we should consider a region with $n_s(K) \cong 1$. This can be satisfied if

$$\begin{aligned} \operatorname{dn}^2(v) (A_1 + A_2 \operatorname{sn}^2(v)) - m^2 \operatorname{sn}^2(v) \operatorname{cn}^2(v) (A_1 + A_2 \operatorname{sn}^2(v)) \\ + 2A_2 \operatorname{sn}^2(v) \operatorname{cn}^2(v) \operatorname{dn}^2(v) = 0 \end{aligned} \quad (26)$$

with the condition

$$\operatorname{sn}(v) \operatorname{cn}(v) \operatorname{dn}(v) (A_1 + A_2 \operatorname{sn}^2(v)) \neq 0. \quad (27)$$

Using some simple relations among Jacobi elliptic functions and introducing a new variable

$$x = \operatorname{sn}^2(v) \quad (28)$$

one gets a cubic equation

$$x^3 + \frac{A_1 m^2 - 4A_2 m^2 - 2A_2}{3A_2 m^2} x^2 + \frac{3A_2 - 2A_1 m^2}{3A_2 m^2} x + \frac{A_1}{3A_2 m^2} = 0 \quad (29)$$

or

$$x^3 + \frac{\rho m^2 - 4m^2 - 2}{3m^2} x^2 + \frac{3 - 2\rho m^2}{3m^2} x + \frac{\rho}{3m^2} = 0, \quad (30)$$

where

$$\rho = \frac{A_1}{A_2} = \frac{\sqrt{\frac{5}{3} - a^2} \sin \frac{\varphi+\pi}{3}}{\sqrt{\frac{5}{3} - a^2} \cos \frac{\varphi+4\pi}{3} + \frac{5+6a}{12}}. \quad (31)$$

This equation can be reduced to

$$y^3 + py + q = 0 \quad (32)$$

where

$$p = \frac{-m^4(\rho^2 + 10\rho + 16) + m^2(4\rho + 11) - 4}{27m^4} \quad (33)$$

$$q = \frac{2m^6(\rho^3 + 15\rho^2 + 92\rho - 64) + 6m^4(49 + 25\rho - 2\rho^2) + 6m^2(43\rho - 64) - 16}{27^2m^6} \quad (34)$$

$$x = y - \left(\frac{\rho m^2 - 4m^2 - 2}{9m^2} \right). \quad (35)$$

The discriminant of Eq. (32) can be written as

$$D = \frac{q^2}{4} + \frac{p^3}{27} \quad (36)$$

and one gets after some tedious algebra

$$\begin{aligned} D(\rho, m) = \frac{1}{27^4 m^{10}} \bigg[& -m^{10} (387\rho^4 + 7376\rho^3 + 29456\rho^2 + 114944\rho + 258048) \\ & + m^8 (243\rho^4 + 6660\rho^3 + 19062\rho^2 + 15000\rho + 116352) \\ & + m^6 (246\rho^4 + 2292\rho^3 - 4716\rho^2 + 3299\rho + 2473) \\ & + m^4 (144\rho^3 + 1512\rho^2 + 10476\rho + 19251) \\ & + m^2 (-1548\rho^3 + 3584\rho^2 + 35190\rho - 26460) \\ & + 48(75 - 39\rho) \bigg]. \end{aligned} \quad (37)$$

We use the Cardano formulae to find a root x_0 . Afterwards we get

$$v_{\pm} = \pm \int_0^{\arcsin \sqrt{x_0}} \frac{d\theta}{\sqrt{1 - m^2 \sin^2 \theta}}. \quad (38)$$

Thus we need $0 \leq x_0 \leq 1$.

It is easy to see that every

$$v_l = v_{\pm} + 2K(m^2)l, \quad l = 0, \pm 1, \pm 2, \dots, \quad (39)$$

satisfies Eq. (26).

If $D(m, \rho) > 0$, we have only one real root

$$x_0 = \sqrt[3]{-\frac{q}{2} + \sqrt{D}} + \sqrt[3]{-\frac{q}{2} - \sqrt{D}} - \left(\frac{\rho m^2 - 4m^2 - 2}{9m^2} \right) \quad (40)$$

where q is given by Eq. (34) and D by Eq. (37). Moreover we need $0 < x_0 < 1$. In order to find a criterion let us transform Eq. (30) via

$$x = \frac{1+w}{1-w}. \quad (41)$$

This transformation maps a unit circle on a complex plane $|x| < 1$ into $\text{Re } w < 0$. One gets

$$\begin{aligned} w^3 + \frac{13m^2 + \rho m^2 - 1 + 3\rho}{7m^2 - 3\rho m^2 + 5 - \rho} w^2 \\ + \frac{5m^2 + 3\rho m^2 - 5 - 3\rho}{7m^2 - 3\rho m^2 + 5 - \rho} w + \frac{\rho + 1 - m^2 - \rho m^2}{7m^2 - 3\rho m^2 + 5 - \rho} = 0. \end{aligned} \quad (42)$$

It is easy to see that the map (41) transforms roots of Eq. (30) into roots of Eq. (42) and real roots into real roots. Especially roots inside the unit circle into roots with negative real part. Thus if Eq. (30) possesses only one real root inside the unit circle then Eq. (42) possesses only one real root with the negative real part. Let w_0 be such a root of Eq. (42) and z and $\bar{z}$ be the remaining roots. One gets

$$\begin{aligned} (w - w_0)(w - z)(w - \bar{z}) \\ = w^3 - (2\text{Re}(z) + w_0)w^2 + (|z|^2 + 2\text{Re}(z)w_0)w - w_0|z|^2 = 0. \end{aligned} \quad (43)$$

Thus we only need

$$-w_0|z|^2 > 0, \quad (44)$$

$$x_0 = \frac{1+w_0}{1-w_0}, \quad (45)$$

which means

$$\frac{\rho + 1 - m^2 - \rho m^2}{7m^2 - 3\rho m^2 + 5 - \rho} > 0. \quad (46)$$

The last condition means also that Eq. (30) has a real root inside the unit circle, i.e. x_0 . Now we only need to satisfy $x_0 > 0$, which simply means

$$\frac{\rho}{3m^2} < 0 \quad (47)$$

or (if $m^2 > 0$)

$$\rho < 0. \quad (48)$$

In this way we get conditions

$$(\rho + 1 - m^2 - \rho m^2)(7m^2 - 3\rho m^2 + 5 - \rho) > 0, \quad (49)$$

$$\rho < 0. \quad (50)$$

From (49–50) we get two solutions:

$$\text{I} \quad 0 > \rho > -1, \quad 1 > m^2 > \frac{5 - \rho}{7 - 3\rho} \quad (51)$$

$$\text{II} \quad \rho < -1, \quad 0 < m^2 < \frac{5 - \rho}{7 - 3\rho}. \quad (52)$$

Simultaneously we have

$$D(\rho, m) > 0. \quad (53)$$

Under these conditions our solution given by Eq. (40) is non-negative with modulus smaller than one.

In this way we have

$$n_s(K) = 1 \quad (54)$$

for

$$\ln K = \frac{v_{\pm} + 2K(m^2)l}{C} + K'_0 \quad (55)$$

$$K = K_0 \exp\left(\frac{v_{\pm} + 2K(m^2)l}{C}\right) \quad (56)$$

where $K'_0 = \ln K_0$.

Moreover we are interested in a deviation from a flat power spectrum. Thus we consider

$$n_s(K) = 1 \pm \frac{dn_s(K)}{d \ln K} \Delta \ln K \quad (57)$$

where $\Delta \ln K \sim 10$ and $\frac{dn_s(K)}{d \ln K}$ is calculated for $\ln K$ given by the formula (54). According to the formula (14) $\frac{dn_s(K)}{d \ln K}$ is proportional to C^2 .

Thus using (25) we get

$$\frac{dn_s(K)}{d \ln K} \Delta \ln K \sim 10 \left(\frac{K(m^2)}{10}\right)^2 \sim 0.1 \quad (58)$$

which gives us rough estimate of a deviation from a flat power spectrum.

For K given by Eq. (56) one gets

$$\begin{aligned} \frac{dn_s(K)}{d \ln K} = 2C^2 & \left[\varepsilon \left(\varepsilon_0 x_0 \sqrt{1 - m^2 x_0} - (1 + x_0(m^2 - 1)) \sqrt{1 - x_0} \right) \right. \\ & \times \frac{\sqrt{1 - m^2 x_0} + \varepsilon \varepsilon_0 m^2 \sqrt{x_0(1 - x_0)}}{\sqrt{x_0(1 - x_0)(1 - m^2 x_0)}} \\ & \left. + \frac{1 + 3m^2 x_0^2 - x_0(2m^2 + 1)}{\rho + x_0} + \frac{4x_0(1 - x_0)(1 - m^2 x_0)}{(\rho + x_0)^2} \right] \end{aligned} \quad (**)$$

where

$$\varepsilon_0^2 = \varepsilon^2 = 1. \quad (***)$$

It is interesting to estimate a range of K . One can consider

$$K_0 \exp\left(\frac{v_+ - K(m^2)}{C}\right) \leq K \leq K_0 \exp\left(\frac{v_+ + K(m^2)}{C}\right) \quad (****)$$

Let us consider Eq. (57). If we find conditions for $\frac{dn_s(K)}{d \ln K}$ to be zero, we get a local flat spectrum (around K given by Eq. (56)). One gets using (28) and (**)

$$\begin{aligned} & \left(\varepsilon_0 x_0 \sqrt{1 - m^2 x_0} - (1 + x_0(m^2 - 1)) \sqrt{1 - x_0} \right) \\ & \times \frac{\sqrt{1 - m^2 x_0} + \varepsilon \varepsilon_0 m^2 \sqrt{x_0(1 - x_0)}}{\sqrt{x_0(1 - x_0)(1 - m^2 x_0)}} \\ & = \frac{x_0(2m^2 + 1) - 1 - 3m^2 x_0^2}{\rho + x_0} + \frac{4x_0(x_0 - 1)(1 - m^2 x_0)}{(\rho + x_0)^2}. \end{aligned} \quad (*****)$$

For $x_0 = x_0(m, \rho)$ (see Eq. (40)) is a known function of m and ρ , the condition (*****) can be considered as a constraint for m and ρ . Together with (51), (52) and (53) it can give an interesting range for parameters m and ρ . In this range the spectrum function (9) looks flat (around K given by Eq. (56)).

Let us consider a formula for an evolution of Higgs' field given by Eq. (3.312) from Ref. [1] corrected in the fifth point of Ref. [2], (14.369):

$$\Phi(t) = \frac{1}{2\alpha_s} \left(1 \pm \sqrt{\frac{5}{1 - e^{-5b(t-t_0)}}} \right) \quad (59)$$

where $b = \frac{e^{n\Psi_1} A}{6r^2 H_0} > 0$,

$$\lim_{t \rightarrow \infty} \Phi(t) = \frac{1}{\alpha_s} \left(\frac{\pm \sqrt{5} + 1}{2} \right).$$

It is easy to see that for

$$t_{\max} - t_0 = \frac{2}{b}, \quad (61)$$

$$\Phi\left(\frac{2}{b} + t_0\right) \simeq \frac{1}{\alpha_s} \left(\frac{1 \pm \sqrt{5}}{2} \right). \quad (62)$$

Thus we can consider from practical point of view that an inflation has been completed for $t = t_0 + \frac{2}{b}$. The inflation starts for

$$t_{\text{initial}} = t_0 + \frac{1}{5b} \ln \frac{5}{4}. \quad (63)$$

Let us calculate $P_R(K)$, a spectral function for this kind of evolution of Higgs' field. One gets from Eqs (7) and (8)

$$P_R(K) = P_0 \frac{\left(\frac{K}{K_0}\right)^{5\bar{\mu}}}{\left(\left(\frac{K}{K_0}\right)^{\bar{\mu}} - 1\right)^3}, \quad (64)$$

$$P_0 = \frac{4H_0^4 \alpha_s^2 n_1}{125\pi^2 b^2} \quad (65)$$

$$K_0 = R_0 H_0 e^{t_0 H_0}. \quad (66)$$

For a spectral index we get

$$\frac{d \ln P_R(K)}{d \ln K} = n_s(K) - 1 = \bar{\mu} \frac{2 \left(\frac{K}{K_0}\right)^{\bar{\mu}} - 5}{\left(\frac{K}{K_0}\right)^{\bar{\mu}} - 1} \quad (67)$$

$$\bar{\mu} = \frac{5b}{H_0}. \quad (68)$$

For the first derivative of n_s

$$\frac{dn_s}{d \ln K} = \frac{8\bar{\mu}^2 \left(\frac{K}{K_0}\right)^{\bar{\mu}}}{\left(\left(\frac{K}{K_0}\right)^{\bar{\mu}} - 1\right)^2}. \quad (69)$$

An interesting question is to find the range of K . Taking

$$t_{\text{initial}}^* = \frac{1}{H_0} \left(\frac{K_{\text{initial}}}{R_0 H_0} \right) = t_0 + \frac{1}{5b} \ln \frac{5}{4} = t_{\text{initial}} \quad (70)$$

one gets

$$K_{\text{initial}} = K_0 \left(\frac{5}{4} \right)^{1/\bar{\mu}} \quad (71)$$

$$t_{\text{max}}^* = \frac{1}{H_0} \left(\frac{K_{\text{max}}}{R_0 H_0} \right) = t_0 + \frac{2}{b} = t_{\text{max}} \quad (72)$$

$$K_{\text{max}} = K_0 e^{2H_0/b} = K_0 e^{10/\bar{\mu}} \quad (73)$$

and finally

$$K_0 \left(\frac{5}{4} \right)^{1/\bar{\mu}} < K < K_0 e^{10/\bar{\mu}}. \quad (74)$$

Let us find $\tilde{K}$ such that

$$n_s(\tilde{K}) = 1. \quad (75)$$

One gets

$$\tilde{K} = K_0 \left(\frac{5}{2} \right)^{1/\bar{\mu}} \quad (76)$$

$$K_0 \left(\frac{5}{4} \right)^{1/\bar{\mu}} < \tilde{K} < K_0 e^{10/\bar{\mu}}. \quad (77)$$

For such a $\tilde{K}$ one gets

$$\frac{dn_s(\tilde{K})}{d \ln K} = 5\bar{\mu}^2 \quad (78)$$

$$n_s(\tilde{K}) \cong 1 \pm 5\bar{\mu}^2 \Delta \ln K. \quad (79)$$

If we take as usual $\Delta \ln K \simeq 10$, we get

$$n_s(K) \cong 1 \pm \frac{1250}{H_0^2} b^2 = 1 \pm \frac{625}{18r^4 H_0^4} e^{2n_{\Psi_1}} A^2. \quad (80)$$

We can try to make it sufficiently closed to 1 taking a big H_0 (see Eq. (2.31) from Ref. [1]).

In this case we have parametrized a spectral function and a spectral index by only three parameters $\bar{\mu}$, P_0 and K_0 , from which only $\bar{\mu}$ matters. Taking K from the range

$$1.2^{1/\bar{\mu}} < \frac{K}{K_0} < 10^{4.3/\bar{\mu}} \quad (81)$$

we can calculate in principle all CMB characteristics (see Ref. [4]) and to fit to observational data (Ref. [5]) (as good as possible) parameters $\bar{\mu}$ (and K_0). This will be a subject of future investigations.

Finally (in order to give a complete set of spectral functions) let us come back to some formulae from Ref. [1] (see Eqs (3.373–376)):

$$P_R(K) = P_0 g \left(\frac{K}{R_0 H_0} \right) \quad (82)$$

$$g(x) = \frac{1}{x^2} \cdot \frac{2}{\left(\text{sd}(u, \frac{1}{2}) + \sqrt{2} \text{cd}(u, \frac{1}{2}) \text{nd}(u, \frac{1}{2}) \right)^2}$$

$$u = \sqrt{2}C_1x + K\left(\frac{1}{2}\right) - C_1\sqrt{2} - \int_0^{\arccos(\sqrt{5}/C_1)} \frac{d\varphi}{\sqrt{1 - \frac{1}{2}\sin^2\varphi}} \quad (83)$$

$$P_0 = \left(\frac{H_0}{2H}\right)^2 \cdot \frac{4\alpha_s^2 n_1}{5C_1^2} \quad (84)$$

$$n_s(K) - 1 = -2 - C_1x \cdot \frac{\sqrt{2}\operatorname{cd}(u, \frac{1}{2}) - \operatorname{sd}^2(u, \frac{1}{2})}{\operatorname{sd}(u, \frac{1}{2}) + \sqrt{2}\operatorname{cd}(u, \frac{1}{2})\operatorname{nd}(u, \frac{1}{2})} \quad (85)$$

$$\begin{aligned} \frac{dn_s(K)}{d\ln K} = & -2C_1x \cdot \frac{\sqrt{2}\operatorname{cd}(u, \frac{1}{2}) - \operatorname{sd}^2(u, \frac{1}{2})}{\operatorname{sd}(u, \frac{1}{2}) + \sqrt{2}\operatorname{cd}(u, \frac{1}{2})\operatorname{nd}(u, \frac{1}{2})} - 2C_1^2x^2\operatorname{sd}^2(u, \frac{1}{2}) \\ & - 2C_1x^2 \cdot \frac{(\sqrt{2}\operatorname{cd}(u, \frac{1}{2}) - \operatorname{sd}^3(u, \frac{1}{2}))^2}{(\operatorname{sd}(u, \frac{1}{2}) + \sqrt{2}\operatorname{cd}(u, \frac{1}{2})\operatorname{nd}(u, \frac{1}{2}))^2} \end{aligned} \quad (86)$$

In this case independent parameters are C_1 and $K_0 = R_0H_0$. The amount of inflation in three cases can be calculated. In the first case (function (17)) it is given by the corrected formulae (3.200–201) from Ref. [1], i.e.

$$\bar{N}_0 = H_0 \sqrt{\frac{2r^2}{5Aae^{n\psi_1}}} \int_{\varphi_2}^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right)\sin^2\theta}} \quad (87)$$

or

$$\bar{N}_0 = \frac{H_0}{\sqrt{5\alpha_s}} \sqrt[4]{\frac{4r^2}{ABe^{n\psi_1}}} \int_{\varphi_2}^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10}\right)\sin^2\theta}} \quad (88)$$

$$\cos^2\varphi_1 = \frac{1}{5+4a}, \quad \cos^2\varphi_2 = \frac{5}{5+4a}. \quad (89)$$

It is interesting to notice that we can calculate $\bar{N}_0$ from Eq. (87) for simplified Weinberg-Salam model (see Ref. [6]) in the nonsymmetric version (see Ref. [2]). One gets

$$\begin{aligned} \bar{N}_0 = & J \cdot \sqrt{\frac{2}{15}} \left(\frac{m_{\tilde{A}}}{m_{\text{pl}}\alpha_s}\right)^{7/2} 2^{-27/4} \\ & \times \frac{\left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)}\right)^{5/4}}{(2\mu^3 + 7\mu^2 + 5\mu + 20)^{7/4} |\zeta|^{1/4} (\mu^2 + 4)^{1/2}} \\ & \times \frac{(1 + \zeta^2)^{1/4} \left(49g^2(\zeta, \mu) + g(\zeta, \mu)(\mu^2 + 4)\sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)}\right)^{1/2}}{\left(\ln(|\zeta| + \sqrt{\zeta^2 + 1}) + 2\zeta^2 + 1\right)^{3/2} ((5\zeta^2 + 4)\ln 2 - 2\mu^2(1 + \zeta^2))^{1/4}} \end{aligned} \quad (87a)$$

where

$$f(\zeta) = \frac{16|\zeta|^3(\zeta^2 + 1)}{3(2\zeta^2 + 1)(1 + \zeta^2)^{5/2}} \left(\zeta^2 E \left(\frac{|\zeta|}{\sqrt{\zeta^2 + 1}} \right) - (2\zeta^2 + 1) K \left(\frac{|\zeta|}{\sqrt{1 + \zeta^2}} \right) \right) + 8 \ln \left(|\zeta| \sqrt{\zeta^2 + 1} \right) + \frac{4(1 + 9\zeta^2 - 8\zeta^4)|\zeta|^3}{3(1 + \zeta^2)^{3/2}} \quad (\text{I})$$

$$g(\zeta, \mu) = -f(\zeta)(\mu^2 + 4) \quad (\text{II})$$

$$h(\zeta, \mu) = |\zeta|(2\mu^3 + 7\mu^2 + 5\mu + 20)(1 + \zeta^2)^{-1} \times ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2)) \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right) \quad (\text{III})$$

$$J = \int_{\varphi_2}^{\varphi_1} \frac{d\theta}{\sqrt{1 - \left(\frac{4a+5}{10} \right) \sin^2 \theta}} \quad (\text{IV})$$

$$a = \left(\frac{m_{\text{pl}}}{m_{\tilde{A}}} \right)^7 2^{35/2} \alpha_s^{16} B^{1/2} \frac{1}{m_{\tilde{A}}} \times \left(\frac{(2\mu^3 + 7\mu^2 + 5\mu + 20)}{(\mu^2 + 4) \left(7g(\zeta, \mu) + \sqrt{49g^2(\zeta, \mu) + 384h(\zeta, \mu)} \right)} \right)^{7/2} \times \frac{(1 + \zeta^2)^{1/2} \left(\ln \left(|\zeta| + \sqrt{\zeta^2 + 1} \right) + 2\zeta^2 + 1 \right)^4}{|\zeta|^{1/2} ((5\zeta^2 + 4) \ln 2 - 2\mu^2(1 + \zeta^2))^{1/2}} \quad (\text{V})$$

and a satisfies the condition (6), which is a condition for an integration constant B , $B > 0$. All of these formulae are subject to conditions

$$\mu > \mu_0 = -3.581552661 \dots \quad (\tilde{R}(\tilde{I}) > 0) \quad (\text{VIa})$$

$$|\zeta| > \zeta_0 = 1.36 \dots \quad (\tilde{P}(\zeta) < 0) \quad (\text{VIb})$$

$$\mu^2 < \frac{(5\zeta^2 + 4) \ln 2}{2(1 + \zeta^2)} \quad (A > 0). \quad (\text{VIc})$$

In the second case considered here the amount of an inflation is infinite (in principle). Moreover, it can be given by a finite formula

$$N = H_0 (t_{\text{end}} - t_{\text{initial}}) = H_0 (t_{\text{max}} - t_{\text{initial}}) = H_0 \left(\frac{2}{b} - \frac{1}{5b} \ln \frac{5}{4} \right) = \frac{H_0}{b} \left(2 - \frac{1}{5} \ln \frac{5}{4} \right) \cong 1.96 \frac{H_0}{b}. \quad (90)$$

In the third case the amount of an inflation is given by the formula (3.364) from Ref. [1]

$$\sqrt{2}C_2 - K \left(\frac{1}{2} \right) = \int_0^{\arccos \left(\frac{1}{2C_1 e^{N_0}} \right)} \frac{d\theta}{\sqrt{1 - \frac{1}{2} \sin^2 \theta}} - \sqrt{2}C_1 e^{N_0}, \quad (91)$$

which is a transcendental equation for N_0 .

All of those $P_R(K)$ functions will be treated by numerical methods from Ref. [4] and afterwards we compare the results with observational data (e.g. Ref. [5]). This will be the first stage of an application of our theory in cosmology. Afterwards we try to estimate intrinsic parameters of our theory and to start a multicomponent inflation.

References

- [1] Kalinowski M. W., *Dynamics of Higgs' field and a quintessence in the nonsymmetric Kaluza–Klein (Jordan–Thiry) theory*, arXiv: hep-th/0306241v1.
- [2] Kalinowski M. W., *Nonsymmetric Fields Theory and its Applications*, World Scientific, Singapore, New Jersey, London, Hong Kong 1990.
Kalinowski M. W., *Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory in a general nonabelian case*, Int. Journal of Theor. Phys. **30**, p. 281 (1991).
Kalinowski M. W., *Nonsymmetric Kaluza–Klein (Jordan–Thiry) Theory in the electromagnetic case*, Int. Journal of Theor. Phys. **31**, p. 611 (1992).
Kalinowski M. W., *Can we get confinement from extra dimensions*, in: Physics of Elementary Interactions (ed. Z. Ajduk, S. Pokorski, A. K. Wróblewski), World Scientific, Singapore, New Jersey, London, Hong Kong 1991.
Kalinowski M. W., *Scalar fields in the nonsymmetric Kaluza–Klein (Jordan–Thiry) theory*, arXiv: hep-th/0307242.
- [3] Liddle A. R., Lyth D. H., *Cosmological Inflation and Large-Scale Structure*, Cambridge Univ. Press, Cambridge 2000.
- [4] www.cmbfast.org
- [5] Bennett C. L. et al., *First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters*, astro-ph/0302209v2.
- [6] Manton N. S., *A new six-dimensional approach to the Weinberg–Salam model*, Nucl. Phys. **B158**, p. 141 (1979).